

On the classical central charge

M. Blagojević and M. Vasilic*

Institute of Physics, P.O.Box 57, 11001 Belgrade, Serbia

Abstract

In the canonical formulation of a classical field theory, symmetry properties are encoded in the Poisson bracket algebra, which may have a central term. Starting from this well understood canonical structure, we derive the related Lagrangian form of the central term.

I. INTRODUCTION

In the canonical description of a classical field theory, the symmetry generators are represented by the phase-space functionals which act on the basic dynamical variables through Poisson brackets (PBs). As a consequence, the commutator algebra of the symmetry transformations is represented by the PB algebra, which may have central extension. If the symmetry group is parametrized by ε^a , and $G = G[\varepsilon]$ is the canonical generator, the PB algebra has the general form [1]

$$\{G[\varepsilon], G[\eta]\} = G[\theta] + K[\varepsilon, \eta], \quad (1.1)$$

where $\theta = \theta(\varepsilon, \eta)$ is defined by the group composition law, and $K[\varepsilon, \eta]$ is a constant phase-space functional—the classical *central term*.

The construction of well defined (finite and differentiable) canonical generators $G[\varepsilon]$ has been thoroughly treated in the Hamiltonian formalism [2–6]. In the first step, one adopts specific asymptotic conditions on the canonical variables, so as to restrict the phase space to the domain on which the canonical generators are well defined. The symmetry group itself is thereby restricted to a subgroup which preserves the asymptotics—the *asymptotic symmetry group*. In the second step, the asymptotic symmetry generators are improved by adding suitable surface terms, whereupon the PB algebra (1.1) naturally emerges.

The adopted asymptotics, defined by the sole requirement of *finiteness* and *differentiability*, is necessary for the algebra (1.1) to make sense, but it does not guarantee the conservation of the related charges—the on-shell values of the asymptotic symmetry generators. In what follows, we shall restrict our interest to finite energy field configurations of isolated physical systems, as their asymptotic behavior ensures the *conservation of the canonical charges*.

*Email addresses: mb@phy.bg.ac.yu, mvasilic@phy.bg.ac.yu

The nature and origin of central terms in algebras associated with physically relevant symmetries is of particular importance for a proper understanding of the quantum features of gravity [7–12]. Central terms in quantum physics do not emerge exclusively as a consequence of the quantum ordering procedure, as in string theory with its Virasoro algebra, but also inherit the underlying classical structure. In the conformal field theory, for instance, in Cardy’s formula for the asymptotic density of states, the dominant contribution involves the classical central term. The same property holds also for the Bekenstein-Hawking entropy of the BTZ black hole. Thus, classical central terms are as physically relevant as their quantum counterparts. The mechanism by which a classical feature (such as classical central term), defined entirely in the realm of classical physics, influences the nature of the quantum theory, is not yet sufficiently clear and deserves further investigation. In addition to this, classical central terms are also related to certain geometric properties of the classical solutions.

Although the structure of central term is well understood at the canonical level, the whole formalism is somewhat indirect: in order to find out the PB algebra with central term, one needs a complete information regarding its Hamiltonian structure (Hamiltonian, canonical generator, etc.). To circumvent this unpractical side of the approach, one would like to find a counterpart of these results within the Lagrangian formalism, which will eventually lead us to the same information in a more direct way.

In the Lagrangian approach to the classical central term, one is effectively faced with the problem of finding the counterpart of the PB algebra (1.1). To this end, we note that this relation can be rewritten as

$$\delta_\eta G[\varepsilon] = G[\theta] + K[\varepsilon, \eta], \quad (1.2)$$

where $\delta_\eta G[\varepsilon]$ is the infinitesimal symmetry transformation of $G[\varepsilon]$ with parameter η . As the on-shell values of $G[\varepsilon]$ represent the canonical conserved charges, it is natural to expect that the transformation properties of the Lagrangian conserved charges may lead us to the Lagrangian counterpart of the canonical central term $K[\varepsilon, \eta]$.

A number of authors in the literature have taken this way to the problem [13–17]. They make use of the fact that continuous symmetries of the classical action lead, via Noether’s theorem [18], to differentially conserved Noether currents. However, Noether current is not defined uniquely, but only up to an exact term, the exterior derivative of the so-called superpotential. In gauge theories the complete conserved current reduces on-shell merely to the superpotential term. Since the superpotential is completely arbitrary, the related Noether charge is essentially undefined. In order to define the Noether charge, one has to find an appropriate criterion which will fix the superpotential. As it turns out, this criterion essentially boils down to the very existence and conservation of Noether charges [15].

Although many important features of the Lagrangian approach have been successfully developed, there are still some open questions; in particular, its equivalence with the Hamiltonian formalism has not been fully understood. The purpose of the present paper is to define central term of the asymptotic symmetry in the Lagrangian context, starting from the related canonical structure (1.1). The approach leads to a very simple but quite general Lagrangian expression for the central term, valid in an arbitrary field theory. In the case of gauge theories, the central term takes the form of a surface integral, as expected. The above procedure represents the first step towards a complete Lagrangian treatment of the asymptotic symmetries, fully equivalent with the canonical structure expressed in (1.1).

The layout of the paper is as follows. In Sect. II, we review some basic aspects of the central term in the canonical formalism, including the canonical origin of this term, and some of its basic properties. In Sect. III, we develop an equivalent Lagrangian description of the central term, leading to an explicit Lagrangian expression for $K[\varepsilon, \eta]$. In Sect. IV we apply these results to gauge theories; in particular, we represent central term in the form of a surface integral. In Sect. V we study several interesting examples: Chern-Simons theory, Liouville theory and general relativity. Section VI is devoted to concluding remarks, while some technical details are relegated to the appendix.

Basic concepts and conventions. Before continuing with the main exposition, we present here a brief account of the basic concepts and conventions used in our paper.

(1) We consider a generic classical field theory defined on an n -dimensional manifold $\mathcal{M}$, conventionally called spacetime. Basic dynamical variables are the fields $\phi = \{\phi^i, i = 1, \dots, m\}$ defined on $\mathcal{M}$; they can be scalars, gauge fields, metric tensor etc. The topology of spacetime $\mathcal{M}$ is assumed to have the form $R^1 \times \Sigma$, where Σ is referred to as the spatial section of $\mathcal{M}$. When the gravitational interaction is neglected, the n -dimensional spacetime has the structure of Minkowski space, but in general, the structure of $\mathcal{M}$ may be quite different. The fields are subject to the field equations derived from an action functional—the integral of a Lagrangian n -form $\mathcal{L}(\phi, \partial\phi)d^n x$, where the Lagrangian function $\mathcal{L}(\phi, \partial\phi)$ is a scalar density.

(2) In addition to the field equations, we impose a set of asymptotic (or boundary) conditions, which restrict the allowed configurations of fields in the asymptotic region. Typically, the asymptotic region is defined as the spatial infinity. In more general situations, the asymptotic region may be different, or even missing (see our second example). Typical asymptotic conditions in gravity correspond to the assumption of flat spacetime structure at spatial infinity, but our approach allows for more general situations, such as anti-de Sitter asymptotics, for instance. In fact, the only limitations on the type of asymptotic behaviour that we need are some quite general requirements on the structure of symmetry generators, as we shall see.

(3) The field equations of our concern are invariant under the action of a continuous group of symmetries U with parameters $\varepsilon = (\varepsilon^a)$. The group U may be a Lie group ($\varepsilon^a = \text{const.}$), but it may also have a more general structure, as in gauge field theories or gravity. Once we adopt the set of asymptotic conditions as an additional element of the dynamical structure of the theory, the original symmetry group U is accordingly modified: it is reduced to the subgroup $U_0 \subseteq U$, consisting of all those transformations that do not change the asymptotic conditions. In this way, the asymptotic conditions on dynamical variables induce the related conditions on the group parameters ε^a .

(4) In the canonical formalism, basic dynamical variables are the functions $\phi(x), \pi(x)$ on Σ , which are the elements of the phase space. By construction, the canonical generator of the symmetry transformations belonging to U_0 is a local functional, $G[\varepsilon, \phi, \pi] = \int_{\Sigma} d^{n-1}x \mathcal{G}(\varepsilon, \partial\varepsilon, \phi, \partial\phi, \pi, \partial\pi)$. When U_0 is a Lie group, we have $G[\varepsilon, \phi, \pi] = \varepsilon^a G_a[\phi, \pi]$. The canonical generator acts on basic dynamical variables via the Poisson bracket operation, which is defined in terms of functional derivatives.

Consider a local functional $F[\phi, \pi] = \int_{\Sigma} d^{n-1}x f(\phi, \pi; \partial\phi, \partial\pi)$, where the functions $\phi(x), \pi(x)$ are defined by certain smoothness properties, asymptotic (or boundary) conditions, and possible additional restrictions [17]. The specific form of these characteristics

depends on the theory we are interested in, and the specific problem we wish to study. The phase space is the space of functions $\Phi : \Sigma \rightarrow R^{2m}$ subject to these additional conditions. A well founded mathematical analysis of the functionals $F[\phi, \pi]$ demands a precise definition of the space Φ , which turns out, in general, to be a rather complicated task (the interested reader may consult [19], for instance).

The functional $F[\phi, \pi]$ is said to have well defined functional derivatives if its variation $\delta F \equiv F[\phi + \delta\phi, \pi + \delta\pi] - F[\phi, \pi]$ can be written in the form

$$\delta F[\phi, \pi] = \int_{\Sigma} d^{n-1}x [A\delta\phi + B\delta\pi] ,$$

where derivatives of $\delta\phi$ and $\delta\pi$ do not appear on the right-hand side [2,4].

Although the canonical generator $G[\varepsilon, \phi, \pi]$ does not satisfy these requirements in general, its form can be improved by adding a suitable surface term, whereupon it becomes differentiable. The procedure depends essentially on the form of asymptotic conditions. In the present discussion, explicit form of the asymptotic conditions is not needed. Instead, we assume that the asymptotic conditions are chosen so as to guarantee differentiability, finiteness and conservation of the improved canonical generators (see section II).

In what follows, we shall use the notation $G[\varepsilon, \phi, \pi]$ for the improved generator, and moreover, the functional dependence on ϕ and π will be often omitted for simplicity, reducing the complete notation $G[\varepsilon, \phi, \pi]$ just to $G[\varepsilon]$.

(5) Asymptotic conditions are an intrinsic part of the canonical formalism, as they define the phase space in which the canonical dynamics takes place. Since these conditions are invariant under the action of the subgroup U_0 of U , it seems natural to take U_0 for the asymptotic symmetry group of the theory. However, in gauge theories and gravity, U_0 contains the set of residual (or pure gauge) transformations, which are defined by the property that their canonical generators weakly vanish. These transformations are nontrivial “inside” $\mathcal{M}$, but act trivially in the asymptotic region. Consequently, they are irrelevant for our discussion of the conserved charges and central terms, both of which are given as some surface integrals. Thus, we are naturally led to introduce the improved definition of the asymptotic symmetry: it is the symmetry defined as the factor group of U_0 with respect to the group of residual symmetries [1].

(6) Our conventions are as follows. In the general analysis, the Latin indices $(i, j, k, \dots)$ are used to count field components, and $(a, b, c, \dots)$ are group indices. The Latin indices $(i, j, k, \dots)$, when used in examples, refer to the local Lorentz frame, the Greek indices $(\mu, \nu, \rho, \dots)$ refer to the coordinate frame, and both run over $0, 1, \dots, n-1$; the Greek indices $(\alpha, \beta, \gamma, \dots)$ are the space indices, and run over $1, 2, \dots, n$; spacetime derivatives are often denoted by comma: $\partial_\mu \phi \equiv \phi_{,\mu}$; totally antisymmetric tensor $\varepsilon^{ij\dots k}$ and the related tensor density $\varepsilon^{\mu\nu\dots\rho}$ are both normalized by $\varepsilon^{01\dots n-1} = +1$; $\eta_{\mu\nu} = (+, -, \dots, -)$ are components of the Minkowski metric, and $\square = \eta^{\mu\nu} \partial_\mu \partial_\nu$.

II. CANONICAL ANALYSIS

In the canonical description of a classical field theory, the basic dynamical variables are the phase-space coordinates (ϕ^i, π_i) . If the theory possesses a symmetry, the equations of motion are invariant under the transformations

$$\delta_\varepsilon \phi^i \equiv \{\phi^i, G[\varepsilon]\}, \quad \delta_\varepsilon \pi_i \equiv \{\pi_i, G[\varepsilon]\}, \quad (2.1)$$

where the canonical generators are phase-space functionals, $G[\varepsilon] \equiv G[\varepsilon, \phi, \pi]$, which act on the phase-space variables through the PB operation. They are finite and differentiable functionals under appropriate asymptotic conditions, which restrict the original symmetry to the so-called asymptotic symmetry [2,4]. The group structure of the symmetry transformations is expressed by the closure of their commutator algebra:

$$[\delta_\eta, \delta_\varepsilon]X = \delta_\theta X, \quad (2.2)$$

where $X = X(\phi, \pi)$, and $(\varepsilon, \eta) \rightarrow \theta(\varepsilon, \eta)$ is the composition law of the group.

The PB algebra. It is clear that adding a constant phase-space functional $c[\varepsilon]$ to the generator $G[\varepsilon]$ will not change its action on the phase space. Therefore, it is not really the phase-space functionals $G[\varepsilon]$ which are in one-to-one correspondence with the symmetry transformations, but their equivalence classes, defined by

$$G[\varepsilon] \sim G[\varepsilon] + c[\varepsilon].$$

This fact has serious consequences on the structure of the underlying canonical symmetry algebra. Indeed, the group structure (2.2) of the symmetry transformations implies

$$\{X, \{G[\varepsilon], G[\eta]\}\} = \{X, G[\theta]\},$$

which guarantees the closure of the PB algebra of the corresponding equivalence classes of generators, while the canonical generators themselves, in general, satisfy the PB algebra with central term,

$$\{G[\varepsilon], G[\eta]\} = G[\theta] + K[\varepsilon, \eta], \quad (2.3)$$

where $K[\varepsilon, \eta]$ is constant over the whole phase space. In particular, if we restrict ourselves to a rigid internal symmetry group, we have $\theta^c = f_{ab}^c \varepsilon^a \eta^b$, and the above algebra takes the simpler form:

$$\{G_a, G_b\} = f_{ab}^c G_c + K_{ab},$$

where we used $G[\varepsilon] = \varepsilon^a G_a$, $K[\varepsilon, \eta] = \varepsilon^a \eta^b K_{ab}$. In what follows, we shall address the important question of how and to what extent the classical central term is related to the structure of a given classical theory.

Asymptotic conditions. Since the canonical generators act via the PB operation, they must be both finite and differentiable functionals [2,4,5]. This property is ensured by a proper construction of $G[\varepsilon]$ and appropriate boundary conditions at spatial infinity, which restrict the allowed values of both the phase-space variables and the symmetry parameters ε . For any two generators $G[\varepsilon]$ and $G[\eta]$, their PB algebra is a realization of the structure of the asymptotic symmetry group, and has the general form (2.3), where $K[\varepsilon, \eta]$ is the canonical central term [1]. Thus, the necessary asymptotic conditions for the algebra (2.3) to make sense are defined by the requirements of

- (i) finiteness and differentiability of the generators.

However, these conditions are not sufficient to ensure the conservation of the related charges. Indeed, every symmetry generator must obey the relation [3,6]

$$\frac{\partial G}{\partial t} + \{G, H_T\} = C_{PFC}, \quad (2.4)$$

where H_T is the total Hamiltonian, and C_{PFC} stands for primary first-class constraints. This relation holds only up to trivial generators, such as constants or surface integrals. Since the temporal change of G is given by the formula

$$\frac{dG}{dt} = \frac{\partial G}{\partial t} + \{G, H_T\},$$

the weak equality $dG/dt \approx 0$ will hold only if the generators are redefined to absorb these constants, and the problematic surface terms are eliminated by the additional asymptotic conditions. In what follows, we shall further restrict the asymptotics by the requirement of

(ii) the conservation of energy.

This will, in addition, imply the conservation of all the canonical charges—the property we need for the consistent Lagrangian treatment of central charges.

Trivial central terms. We have already seen that additive constants in the canonical generators do not influence the symmetry transformation law (2.1). However, as seen from (2.3), such a change of $G[\varepsilon]$ induces a change of the central term:

$$G[\varepsilon] \rightarrow G[\varepsilon] + c[\varepsilon] \quad \Rightarrow \quad K[\varepsilon, \eta] \rightarrow K[\varepsilon, \eta] - c[\theta]. \quad (2.5)$$

Conversely, if a central term has the form $K[\theta(\varepsilon, \eta)]$, it is always possible to redefine the generators according to $G[\varepsilon] \rightarrow G[\varepsilon] + K[\varepsilon]$ so that the new central term vanishes.

- (1) Any central term which can be eliminated by the simple redefinition of symmetry generators $G[\varepsilon] \rightarrow G[\varepsilon] + c[\varepsilon]$, is considered trivial.

We shall often ignore trivial central terms, as they can always be transformed to zero.

There are two particular cases in which one can immediately recognize trivial central terms [1,15]. The first is defined by the existence of at least one phase-space point invariant under the action of the symmetry group. Indeed, if ϕ^i and π_i have vanishing PBs with all the generators $G[\varepsilon]$ at some point $(\bar{\phi}, \bar{\pi})$, then $\partial G[\varepsilon]/\partial \phi^i$ and $\partial G[\varepsilon]/\partial \pi_i$ vanish at $(\bar{\phi}, \bar{\pi})$. As a consequence, the PB $\{G[\varepsilon], G[\eta]\}$ also vanishes, and the central term takes the form $K[\varepsilon, \eta] = -G[\theta, \bar{\phi}, \bar{\pi}]$. Therefore,

- (2a) if there exists a phase-space point invariant under the action of the symmetry group, the corresponding central term is necessarily trivial.

The second case refers to symmetry groups whose group structure does not allow for non-trivial central extensions. Indeed, applying the Jacobi identity to (2.3), we obtain relations which restrict the values of $K[\varepsilon, \eta]$:

$$K[\theta(\varepsilon, \eta), \tau] + \text{cyclic}(\varepsilon, \eta, \tau) = 0.$$

As a consequence, it may happen that this constraint has no non-trivial solutions for $K[\varepsilon, \eta]$. Such an example is described by the following theorem:

- (2b) If the algebra of symmetry generators is a semi-simple Lie algebra, the central term is necessarily trivial.

The proof is based on the observation that the constraint $f_{ab}{}^d K_{dc} + \text{cyclic}(a, b, c) = 0$ has no nontrivial solutions for K_{ab} if the structure constants $f_{ab}{}^c$ define a semi-simple Lie algebra.

Canonical equivalence. Given a Hamiltonian classical theory, one can always redefine phase-space variables to obtain canonically equivalent theory. This is done by means of canonical transformations,

$$\phi \rightarrow \phi' = \phi'(\phi, \pi), \quad \pi \rightarrow \pi' = \pi'(\phi, \pi),$$

defined by demanding the invariance of the PB structure:

$$\{A', B'\}' = \{A, B\}$$

for every $A' \equiv A'(\phi', \pi') = A(\phi, \pi)$, and similarly for B' . The symmetry generators of the canonically equivalent theory are simply $G'[\varepsilon, \phi', \pi'] = G[\varepsilon, \phi, \pi]$, and therefore,

$$\{G'[\varepsilon], G'[\eta]\}' = \{G[\varepsilon], G[\eta]\} = G[\theta] + K[\varepsilon, \eta] = G'[\theta] + K[\varepsilon, \eta]. \quad (2.6)$$

As we can see, the central term remains unchanged:

- (3) Canonically equivalent Hamiltonian theories have identical central terms.

Going over to the Lagrangian formalism, one can make use of the above theorem to prove two similar theorems concerning the equality of central terms in classically equivalent Lagrangian theories. We note that adding a pure divergence term to the Lagrangian leads to canonically equivalent formulation of the theory. The above theorem then implies:

- (4a) A pure divergence term in the Lagrangian does not influence the central term.

Another way of obtaining an equivalent Lagrangian theory is to make a different choice of variables. For example, the so-called contact transformations $\phi \rightarrow \phi' = \phi'(\phi, t)$ are known to produce canonically equivalent Hamiltonian theories. Indeed, since the canonical momenta of the transformed Lagrangian $\mathcal{L}'(\phi', \partial\phi') \equiv \mathcal{L}(\phi, \partial\phi)$ are $\pi'_i \equiv \partial\mathcal{L}'/\partial\dot{\phi}'^i = (\partial\phi^k/\partial\phi'^i)\pi_k$, it is not difficult to verify that the transformation $(\phi, \pi) \rightarrow (\phi', \pi')$ is canonical. As canonical transformations do not change the central term, the following theorem is proved:

- (4b) Contact transformations of the Lagrangian variables induce only trivial changes of the central term.

In particular, if we make the change of variables $\phi = \bar{\phi} + \varphi$, where $\bar{\phi}$ is a fixed field configuration (usually a solution of the field equations), the related change of the central term is trivial.

At the end of this section, let us mention one more way of changing Lagrangian without changing the classical field equations: multiplying the Lagrangian by a constant, $\mathcal{L} \rightarrow \alpha \mathcal{L}$. As we shall see in the next section, the central term is thereby changed as $K \rightarrow \alpha K$. Since the corresponding quantum theories are not equivalent (the functional integrals defined by $\mathcal{L}$ and $\alpha \mathcal{L}$ are different), we can say that classical central charges are not purely classical objects after all.

III. CENTRAL CHARGE IN THE LAGRANGIAN FORMALISM

In this section we express the canonical central term through the Lagrangian coefficients, and establish a basic relation between the canonical symmetry generator and the Noether charge.

A. Canonical central charge

We consider a field theory defined by the action $I = \int d^n x \mathcal{L}(\phi, \partial\phi)$. The action is invariant (up to a boundary term) under the symmetry transformations

$$\delta_\varepsilon \phi^i = R^i(\varepsilon, \partial\varepsilon, \phi, \partial\phi), \quad (3.1)$$

which constitute a group U with parameters ε^a . Moreover, we assume that the adopted asymptotic conditions are invariant under a subgroup U_0 of U , which essentially coincides with the asymptotic symmetry group.

After introducing the notation $\varphi^i = \phi^i - \bar{\phi}^i$, where $\bar{\phi}$ is a fixed configuration of fields, the Lagrangian function can be expressed as a series in φ and $\partial\varphi$:

$$\begin{aligned} \mathcal{L} = & V + V_i \varphi^i + V_i^\mu \partial_\mu \varphi^i \\ & + V_{ij} \varphi^i \varphi^j + V_{ij}^\mu \partial_\mu \varphi^i \varphi^j + V_{ij}^{\mu\nu} \partial_\mu \varphi^i \partial_\nu \varphi^j + \mathcal{O}_3, \end{aligned} \quad (3.2)$$

where $\mathcal{O}_n$ denotes a term of the n -th power in φ and/or $\partial\varphi$, and the coefficients $(V, V_i, \dots)$ are functions of $\bar{\phi}$. Let us now introduce the generalized momentum variables

$$\begin{aligned} \Pi_i^\mu &= \frac{\partial \mathcal{L}}{\partial \partial_\mu \phi^i} = V_i^\mu + V_{ij}^\mu \varphi^j + 2V_{ij}^{\mu\nu} \partial_\nu \varphi^j + \mathcal{O}_2, \\ \pi_i^\mu &= \Pi_i^\mu - \bar{\Pi}_i^\mu = V_{ij}^\mu \varphi^j + 2V_{ij}^{\mu\nu} \partial_\nu \varphi^j + \mathcal{O}_2, \end{aligned}$$

where $\bar{\Pi}_i^\mu \equiv \Pi_i^\mu(\phi = \bar{\phi})$. The variables $(\varphi^i, \pi_i \equiv \pi_i^0)$ are canonical, as they are obtained from the original variables $(\phi^i, \Pi_i \equiv \Pi_i^0)$ by the canonical transformation, but $\pi_i \neq \partial \mathcal{L} / \partial \dot{\varphi}^i$. The action of the symmetry transformations (3.1) on (φ^i, π_i) is given by

$$\begin{aligned} \delta_\varepsilon \varphi^i &\equiv \bar{R}^i(\varepsilon) + \mathcal{O}_1, \\ \delta_\varepsilon \pi_i &= V_{ij}^0 \bar{R}^j(\varepsilon) + 2V_{ij}^{0\nu} \partial_\nu \bar{R}^j(\varepsilon) + \mathcal{O}_1, \end{aligned} \quad (3.3a)$$

where $\bar{R}^i(\varepsilon)$ is short for $R^i(\varepsilon, \partial\varepsilon, \bar{\phi}, \partial\bar{\phi})$.

We now wish to construct the canonical generator of the transformations (3.3a). To this end, note that both equations (3.3a) and those which define π_i can be compared with the corresponding canonical relations only on shell, using the Hamiltonian dynamical equations for $\dot{\varphi}^i$. This is, in fact, true in general: the Lagrangian and Hamiltonian approaches can be compared only on shell. However, for arbitrary $\bar{\phi}$ the Lagrangian field equations take the form $V^i - \partial^\mu V_\mu^i + \mathcal{O}_1(\varphi, \partial\varphi) = 0$, so that, on shell, $\mathcal{O}_0$ and $\mathcal{O}_1$ terms are not independent. To avoid this conflicting situation, we assume that $\bar{\phi}$ is a *solution of the field equations*, whereby the clean separation between $\mathcal{O}_0$ and $\mathcal{O}_1$ terms is maintained on shell, too. Then, the canonical generator of the transformations (3.3a) can be written in the form

$$\begin{aligned}
G[\varepsilon] &= \int d^{n-1}x \mathcal{G}(\varepsilon), \\
\mathcal{G}(\varepsilon) &= \bar{R}^i(\varepsilon)\pi_i - \varphi^i \left[V_{ij}^0 \bar{R}^j(\varepsilon) + 2V_{ij}^{0\nu} \partial_\nu \bar{R}^j(\varepsilon) \right] + \mathcal{O}_2(\varphi, \pi),
\end{aligned} \tag{3.3b}$$

Indeed, the canonical analogs of the transformation laws (3.3a), derived from (3.3b), contain $\mathcal{O}_1(\varphi, \pi)$ terms. Since π is linear in $(\varphi, \partial\varphi)$, we have $\mathcal{O}_1(\varphi, \pi) \approx \mathcal{O}_1(\varphi, \partial\varphi)$ (on shell), and consequently, (3.3a) follows consistently from (3.3b).

We see that the canonical generator (3.3b) satisfies the needed differentiability condition in the lowest order, including only $\mathcal{O}_1(\varphi, \pi)$ terms. With this accuracy, the canonical algebra (2.3) takes the form

$$\{G[\varepsilon], G[\eta]\} = K[\varepsilon, \eta] + \mathcal{O}_1(\varphi, \pi), \tag{3.4a}$$

where

$$\begin{aligned}
K[\varepsilon, \eta] &= \int d^{n-1}x \mathcal{K}(\varepsilon, \eta), \\
\mathcal{K}(\varepsilon, \eta) &= \bar{R}^i(\varepsilon) \left[V_{ij}^0 \bar{R}^j(\eta) + 2V_{ij}^{0\nu} \partial_\nu \bar{R}^j(\eta) \right] - (\varepsilon \leftrightarrow \eta)
\end{aligned} \tag{3.4b}$$

is the classical central charge. It is identified as the zero order term on the right-hand side of (3.4a). The identification is unique since $\bar{\phi}$ is a solution of the field equations.

There is an ambiguity in the derivation of the formula (3.4), which stems from the fact that the coefficients in the $\mathcal{O}_1(\varphi, \pi)$ part of the canonical generator (3.3b) are not defined uniquely. Indeed, one can change any of these coefficients by adding a term proportional to an undetermined Hamiltonian multiplier $v(t)$ (suitably restricted by the adopted asymptotic conditions), since such a term is of the $\mathcal{O}_0(\varphi, \pi)$ type, while its Lagrangian counterpart is some undetermined velocity, thus of the $\mathcal{O}_1$ type. Therefore, the action of such a modified canonical generator will produce the same $\mathcal{O}_0$ terms as in (3.3a). Can one resolve this ambiguity without examining higher order terms in the transformation law?

Let us recall that our choice of the asymptotic conditions includes the requirement of the conservation of energy. This means that the improved total Hamiltonian H_T , which contains an additive surface term S , should satisfy

$$\frac{dH_T}{dt} = \frac{\partial H_T}{\partial t} = C_{PFC} + \frac{\partial S}{\partial t} \approx 0.$$

Thus, $\partial S/\partial t \approx 0$, i.e. the surface integral S has no explicit time dependence, and therefore, no $v(t)$ terms. Using this property of H_T in the derivation of equation (2.4), one can show that there are no surface terms on the right-hand side of that equation, and consequently, all the symmetry generators are also conserved, $dG/dt \approx 0$. Now, combining (1.1) and (2.4) one finds that the central charge is time independent:

$$\frac{\partial}{\partial t} K[\varepsilon, \eta] = 0,$$

for every ε and η , and independently, for every choice of $v(t)$. This implies that there can be no $v(t)$ terms in the central charge $K[\varepsilon, \eta]$. Consequently, our formula (3.4) gives a correct Lagrangian expression for the canonically defined classical central charge.

The general theorem (4b) of the preceding section states that the choice of $\bar{\phi}$ is inessential for the value of the central charge, as a consequence of the canonical invariance of the theory. On the other hand, one can see from our result (3.4) that $K[\varepsilon, \eta]$ explicitly depends on $\bar{\phi}$. This apparent contradiction is resolved by noting that our choice of $\bar{\phi}$ is naturally adjusted to the asymptotic properties of the theory, in the sense that

- (a) $\bar{\phi}$ is a solution of the field equations, and
- (b) it satisfies the adopted asymptotic conditions.

As a consequence, the original *canonical invariance is broken*, and $\bar{\phi}$ cannot be changed arbitrarily any more. Still, there remains the freedom to replace $\bar{\phi}$ by another field configuration satisfying (a) and (b), which produces only a trivial modification of the central charge. The dependence of $K[\varepsilon, \eta]$ on $\bar{\phi}$ is quite natural, as our choice of $\bar{\phi}$ embodies the basic structure of the asymptotic symmetry.

For further analysis and comparison with the Lagrangian formalism, it is useful to define the “canonical current”

$$\mathcal{G}^\mu(\varepsilon) = \varphi^i \left[V_{ij}^\mu \bar{R}^j(\varepsilon) + 2V_{ij}^{\mu\nu} \partial_\nu \bar{R}^j(\varepsilon) \right] - \bar{R}^i(\varepsilon) \left[V_{ij}^\mu \varphi^j + 2V_{ij}^{\mu\nu} \partial_\nu \varphi^j \right] + \mathcal{O}_2, \quad (3.5a)$$

as a generalization of the canonical generator density $\mathcal{G}$ (up to a sign), and the “central current”

$$\mathcal{K}^\mu(\varepsilon, \eta) \equiv \bar{R}^i(\varepsilon) \left[V_{ij}^\mu \bar{R}^j(\eta) + 2V_{ij}^{\mu\nu} \partial_\nu \bar{R}^j(\eta) \right] - (\varepsilon \leftrightarrow \eta), \quad (3.5b)$$

as a generalization of the central charge density $\mathcal{K}$. These two currents satisfy the relation

$$\delta_\varepsilon \mathcal{G}^\mu(\eta) = \mathcal{K}^\mu(\varepsilon, \eta) + \mathcal{O}_1, \quad (3.6)$$

which is the “current” version of the canonical algebra (3.4a).

B. Noether current

The symmetry transformation (3.1) leaves the action invariant, up to a boundary term: $\delta_\varepsilon I = \int_{\mathcal{M}} dK$, where $K = K(\varepsilon, \partial\varepsilon, \phi, \partial\phi)$ is an $(n-1)$ -form on $\mathcal{M}$. Since $dK = \partial_\mu k^\mu d^n x$, where $k^\mu \equiv \varepsilon^{\mu\mu_1 \dots \mu_{n-1}} K_{\mu_1 \dots \mu_{n-1}} / (n-1)!$, this identity can be written as

$$\delta_\varepsilon \mathcal{L} = \partial_\mu k^\mu. \quad (3.7a)$$

The coefficients k^μ are the components of the vector-valued n -form $k^\mu \partial_\mu \otimes d^n x$. Equation (3.7a) implies

$$F_i \delta_\varepsilon \phi^i = \partial_\mu N^\mu. \quad (3.7b)$$

Here, $F_i = \delta \mathcal{L} / \delta \phi^i$ are the field equations, and N^μ is the (conserved) Noether current:

$$N^\mu = k^\mu - \frac{\partial \mathcal{L}}{\partial \partial_\mu \phi^i} \delta_\varepsilon \phi^i \equiv N^\mu(\varepsilon), \quad (3.7c)$$

where, as usual, $N^\mu(\varepsilon)$ is short for $N^\mu(\varepsilon, \partial\varepsilon, \phi, \partial\phi)$. When U_0 is a Lie group, the Noether current takes the form $N^\mu(\varepsilon) = \varepsilon_a N^{a\mu}(\phi, \partial\phi)$. One should note that N^μ is defined only up to a trivial term $\partial_\rho N^{\mu\rho}$, with $N^{\mu\rho} = -N^{\rho\mu}$. We shall show that there is a suitable choice of the superpotential $N^{\mu\rho}$, such that $\mathcal{G}^\mu = N^\mu + \mathcal{O}_2$, which identifies N^μ as the Lagrangian equivalent of $\mathcal{G}^\mu$.

As $\bar{\phi}$ is a solution of the field equations, it holds $V_i = \partial_\mu V_i^\mu$, which implies

$$\begin{aligned}\mathcal{L} &= V + \partial_\mu(V_i^\mu \varphi^i) + \mathcal{L}_2 \\ \mathcal{L}_2 &\equiv V_{ij} \varphi^i \varphi^j + V_{ij}^\mu \partial_\mu \varphi^i \varphi^j + V_{ij}^{\mu\nu} \partial_\mu \varphi^i \partial_\nu \varphi^j + \mathcal{O}_3.\end{aligned}$$

The current k^μ is now given by

$$k^\mu = V_j^\mu R^j + \ell^\mu, \quad \delta_\varepsilon \mathcal{L}_2 = \partial_\mu \ell^\mu.$$

Looking for ℓ^μ in the form

$$\ell^\mu = \alpha^\mu + \alpha_j^\mu \varphi^j + \alpha_j^{\mu\nu} \partial_\nu \varphi^j + \mathcal{O}_2,$$

and using the arbitrariness of φ and $\partial\varphi$, we find that the coefficients satisfy the following set of cascade equations:

$$\begin{aligned}\partial_\mu \alpha^\mu &= 0, \\ \partial_\mu \alpha_j^\mu &= 2V_{ij} \bar{R}^i + V_{ij}^\mu \partial_\mu \bar{R}^i, \\ \alpha_j^\mu + \partial_\rho \alpha_j^{\rho\mu} &= V_{ji}^\mu \bar{R}^i + 2V_{ji}^{\mu\rho} \partial_\rho \bar{R}^i, \\ \alpha_j^{\mu\rho} + \alpha_j^{\rho\mu} &= 0.\end{aligned}\tag{3.8}$$

The first equation implies $\alpha^\mu = \partial_\rho \alpha^{\mu\rho}$, with $\alpha^{\mu\rho} = -\alpha^{\rho\mu}$, so that

$$\begin{aligned}\ell^\mu &= (\alpha_j^\mu - \partial_\rho \alpha_j^{\rho\mu}) \varphi^j + \partial_\rho \ell^{\mu\rho} + \mathcal{O}_2, \\ \ell^{\mu\rho} &\equiv (\alpha^{\mu\rho} + \alpha_j^{\mu\rho} \varphi^j).\end{aligned}$$

Disregarding the trivial term $\partial_\rho \ell^{\mu\rho}$ leads to

$$k^\mu = V_j^\mu R^j + (V_{ji}^\mu \bar{R}^i + 2V_{ji}^{\mu\rho} \partial_\rho \bar{R}^i) \varphi^j + \mathcal{O}_2.$$

Finally, combining this result with the definition (3.7c) of the Noether current, we obtain

$$\begin{aligned}N^\mu &= k^\mu - R^i (V_i^\mu + V_{ij}^\mu \varphi^j + 2V_{ij}^{\mu\nu} \partial_\nu \varphi^j) + \mathcal{O}_2 \\ &= \varphi^j (V_{ji}^\mu \bar{R}^i + 2V_{ji}^{\mu\rho} \partial_\rho \bar{R}^i) - \bar{R}^i (V_{ij}^\mu \varphi^j + 2V_{ij}^{\mu\nu} \partial_\nu \varphi^j) + \mathcal{O}_2 \\ &= \mathcal{G}^\mu + \mathcal{O}_2.\end{aligned}\tag{3.9}$$

Let us stress that this equality holds only if we choose $\partial_\rho \ell^{\mu\rho} = 0$. Accordingly, equation (3.9) can be understood as a condition which defines the choice of the superpotential $N^{\mu\rho}$.

The conditions (3.8) on α_i^μ imply the identity

$$2V_{ij} \bar{R}^i + V_{ij}^\mu \partial_\mu \bar{R}^i = \partial_\mu (V_{ji}^\mu \bar{R}^i + 2V_{ji}^{\mu\rho} \partial_\rho \bar{R}^i), \tag{3.10}$$

directly related to the invariance condition (3.7). Using this identity, we easily derive the relation

$$\partial_\mu \mathcal{K}^\mu(\varepsilon, \eta) = 0,$$

which implies $\mathcal{K}^\mu = \partial_\rho \mathcal{K}^{\mu\rho}$, with $\mathcal{K}^{\mu\rho} = -\mathcal{K}^{\rho\mu}$ (Poincaré lemma). Consequently, the central term can be represented as a surface integral:

$$K[\varepsilon, \eta] = \int d^{n-1}x \mathcal{K}^0(\varepsilon, \eta) = \int dS_\beta \mathcal{K}^{0\beta}(\varepsilon, \eta). \quad (3.11)$$

The conditions of the Poincaré lemma ensure that this result holds true in topologically trivial spaces, while in general, central charges may fail to be represented by surface integrals.

C. Algebra of charges

As we have seen in the preceding subsection, the Noether current N^μ can be chosen to equal the canonical current $\mathcal{G}^\mu$ in the lowest order. This allows us to rewrite (3.6) in a pure Lagrangian form:

$$\delta_\varepsilon N^\mu(\eta) = \mathcal{K}^\mu(\varepsilon, \eta) + \mathcal{O}_1. \quad (3.12)$$

It seems that this relation can be used as the Lagrangian definition of the central charge. Here, the Noether current is identified with the canonical current $\mathcal{G}^\mu$. Can we similarly identify the Noether conserved charges $Q[\varepsilon]$ with the canonical generators $G[\varepsilon]$? A short analysis shows that the answer is affirmative. Namely, it is well known that the well defined canonical generators $G[\varepsilon]$ are also conserved quantities of the theory. If we define the Lagrangian charges

$$Q \equiv G\left[\phi, \Pi = \frac{\partial \mathcal{L}}{\partial \dot{\phi}}\right]_{\phi=\bar{\phi}+\varphi}, \quad (3.13)$$

they will certainly be conserved as the on-shell values of the conserved generators. Moreover, the transformation properties of the charges $Q[\varepsilon]$ will match those of $G[\varepsilon]$:

$$\delta_\eta Q[\varepsilon] \approx Q[\theta] + K[\varepsilon, \eta]. \quad (3.14a)$$

This is a simple consequence of the fact that symmetry transformations do not change the field equations. The weak equality in (3.14a) means the equality up to terms proportional to the equations of motion. Notice, however, there is no physical distinction between $Q[\varepsilon]$ and $Q[\varepsilon] + \mathcal{O}(F)$ since their on-shell values are the same. It is, therefore, natural to work with the equivalence classes of charges $\langle Q \rangle$, defined by

$$\langle Q_1 \rangle = \langle Q_2 \rangle \quad \Leftrightarrow \quad Q_1 \approx Q_2.$$

In terms of the equivalence classes, the algebra of charges is given by the strong equality

$$\delta_\eta \langle Q[\varepsilon] \rangle = \langle Q[\theta] \rangle + K[\varepsilon, \eta]. \quad (3.14b)$$

The algebra of charges (3.14) yields a Lagrangian definition of the central charge. Notice, however, that neither the Noether current $N^\mu(\varepsilon)$ nor the charge $Q[\varepsilon]$ are uniquely defined.

The closure of the algebra (3.14) is obtained only if the existing arbitrariness is fixed in a definite, precise way. We have seen that this can be achieved by identifying Lagrangian charges with the on-shell values of canonical generators, and choosing the asymptotic conditions which make canonical generators well defined and ensure the conservation of energy. If we do not want to refer to the Hamiltonian theory, the only way to find the proper Noether superpotential and boundary conditions is to demand finiteness and conservation of Noether charges [14].

IV. CENTRAL CHARGE IN GAUGE THEORIES

Gauge theories are by far the most interesting field theories from the physical point of view. In the case of gauge symmetries, we derive an explicit representation of the central term as a surface integral.

A. Noether identities

Consider a gauge theory with the transformation law

$$\delta_\varepsilon \phi^i = R_a^i \varepsilon^a + R_a^{i\mu} \partial_\mu \varepsilon^a. \quad (4.1)$$

In this case, the invariance condition (3.7b) can be written in the form

$$\left[F_i R_a^i - \partial_\mu (F_i R_a^{i\mu}) \right] \varepsilon^a = \partial_\mu J^\mu, \quad J^\mu \equiv N^\mu - F_i R_a^{i\mu} \varepsilon^a. \quad (4.2)$$

Let Ω be a region in spacetime, and let the gauge parameters $\varepsilon^a = \varepsilon^a(x)$ satisfy the conditions: a) ε^a are arbitrary in Ω , and b) ε^a and their derivatives vanish on $\partial\Omega$. Then, the integration of equation (4.2) over Ω leads to the following identities:

$$F_i R_a^i - \partial_\mu (F_i R_a^{i\mu}) = 0, \quad (4.3a)$$

$$\partial_\mu J^\mu = 0. \quad (4.3b)$$

The first identity is known as the Noether identity, and the second one implies (Poincaré lemma) that

$$J^\mu = \partial_\rho N^{\mu\rho}, \quad N^{\mu\rho} = -N^{\rho\mu}. \quad (4.3c)$$

As a consequence, we find that Noether current has the form

$$N^\mu = F_i R_a^{i\mu} \varepsilon^a + \partial_\rho N^{\mu\rho}. \quad (4.4)$$

One should stress that, here, $N^{\mu\rho}$ is a completely arbitrary superpotential. Its value is fixed by the extra condition (3.9), which relates Noether and canonical structures of the theory.

Having in mind our goal, to find the Lagrangian structure of the canonical central term, it is useful to expand the Noether identity in terms of φ and $\partial\varphi$. Starting from the Lagrangian (3.2) where $\phi = \bar{\phi}$ is a solution of the field equations, $V_i - \partial_\mu V_i^\mu = 0$, we obtain

$$\begin{aligned} F_i &= (2V_{ij} - \partial_\rho V_{ij}^\rho) \varphi^j + 2(V_{ji}^\nu - \partial_\rho V_{ji}^{\nu\rho}) \partial_\nu \varphi^j - 2V_{ij}^{\mu\nu} \partial_\mu \partial_\nu \varphi^j + \mathcal{O}_2 \\ &\equiv f_{ij} \varphi^j + f_{ij}^\nu \partial_\nu \varphi^j + f_{ij}^{\mu\nu} \partial_\mu \partial_\nu \varphi^j + \mathcal{O}_2. \end{aligned} \quad (4.5)$$

Using this result, the Noether identity (4.3a) takes the form given in Appendix A, which will be useful in further considerations.

B. Central term as a surface integral

Starting from the equality $N^\mu = \mathcal{G}^\mu + \mathcal{O}_2$, equation (3.9), it follows that $\mathcal{G}^\mu$ has the Noether form (4.4):

$$\mathcal{G}^\mu(\varepsilon) = F_i \bar{R}_a^{i\mu} \varepsilon^a + \partial_\rho N^{\mu\rho} + \mathcal{O}_2. \quad (4.6a)$$

Using the expression (3.5a) for $\mathcal{G}^\mu(\varepsilon)$, one finds that $N^{\mu\lambda}$ is given by (Appendix B):

$$\begin{aligned} N^{\mu\lambda} = & \left\{ \left[(V_{ji}^\mu - \partial_\rho V_{ji}^{\mu\rho}) \varphi^j - 2V_{ji}^{(\mu\rho)} \partial_\rho \varphi^j \right] \bar{R}_a^{i\lambda} \varepsilon^a \right. \\ & \left. + V_{ji}^{\mu\lambda} \varphi^j \bar{R}^i + \frac{2}{3} \partial_\rho (V_{ji}^{\mu\rho} \varphi^j \bar{R}_a^{i\lambda} \varepsilon^a) \right\} - (\mu \leftrightarrow \lambda) + \mathcal{O}_2. \end{aligned} \quad (4.6b)$$

Now, the relation $\delta_\varepsilon \mathcal{G}^\mu(\eta) = \mathcal{K}^\mu(\varepsilon, \eta) + \mathcal{O}_1$ and the above expression for $\mathcal{G}^\mu$ imply

$$\mathcal{K}^\mu(\varepsilon, \eta) = \partial_\lambda \left[\delta_\varepsilon N^{\mu\lambda}(\eta) \right]_{\varphi=0}. \quad (4.7a)$$

Consequently, the central term is expressed as the following surface integral:

$$K[\varepsilon, \eta] = \int d^{n-1}x \mathcal{K}^0(\varepsilon, \eta) = \int dS_\beta \left[\delta_\varepsilon N^{0\beta}(\eta) \right]_{\varphi=0}. \quad (4.7b)$$

V. EXAMPLES

A. Chern-Simons theory

As our first example, we consider a Chern-Simons gauge theory on a 3d spacetime $\mathcal{M}$ with the topology $\mathcal{M} = R \times \Sigma$, where R is interpreted as time, and Σ is a spatial manifold whose boundary is topologically a circle located at infinity. For the gauge group $SL(2, R)$, the theory is defined by the Lagrangian

$$\mathcal{L}(A, \partial A) = \kappa \varepsilon^{\mu\nu\rho} \left(A_\mu^a \partial_\nu A_{a\rho} + \frac{1}{3} \varepsilon_{abc} A_\mu^a A_\nu^b A_\rho^c \right). \quad (5.1a)$$

The action is invariant under the infinitesimal gauge transformations

$$\delta_\tau A_\mu = \nabla_\mu \tau \equiv \partial_\mu \tau + [A_\mu, \tau], \quad (5.1b)$$

where $A_\mu = A_\mu^a T_a$, $\tau = \tau^a T_a$, and T_a is the basis of the Lie algebra $sl(2, R)$, defined by $[T_a, T_b] = \varepsilon_{ab}^c T_c$. The field equations have the form

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu] = 0. \quad (5.1c)$$

Every solution of the field equations is a pure gauge: $A_\mu = g^{-1} \partial_\mu g$, where g is in $SL(2, R)$.

We choose the asymptotic conditions which, in polar coordinates $x^\mu = (t, r, \varphi)$, read:

$$A_+ \equiv \frac{1}{2}(A_0 + A_2) = \mathcal{O}_1, \quad (5.2)$$

and $\mathcal{O}_1$ is a term that tends to zero as $1/r$ or faster when $r \rightarrow \infty$. The conditions (5.2) imply that gauge parameters are independent of $x^+ = x^0 + x^2$ in the asymptotic region. In what follows, we shall often refine the asymptotics by the assumption that field equations decrease arbitrarily fast in spatial infinity, since no solution of the field equations is thereby lost. Using this principle, we find that $\partial_+ A_\mu = \mathcal{O}_1$, whereupon the asymptotic dependence on x^+ is completely eliminated from the theory.

In what follows, we shall analyze two typical situations belonging to the class of asymptotic conditions defined above. As before, our fields are expanded around a classical solution,

$$A_\mu = \bar{A}_\mu + \mathcal{A}_\mu, \quad \delta_\tau \mathcal{A}_\mu = \bar{\nabla}_\mu \tau + [\mathcal{A}_\mu, \tau], \quad (5.3)$$

where $\bar{A}_\mu$ satisfies (5.2).

1. We begin by noting that gauge transformations in (5.3) are defined in terms of the gauge parameters τ^a . The related commutator algebra has the form $[\delta_\lambda, \delta_\tau] \mathcal{A}_\mu^a = \delta_\theta \mathcal{A}_\mu^a$, where $\theta^a = \varepsilon^a_{bc} \lambda^b \tau^c$. The central term is defined by the relation (3.4), where

$$\bar{R}^i(\tau) V_{ij}^0 \bar{R}^j(\lambda) - (\tau \leftrightarrow \lambda) = 2\kappa \varepsilon^{0\alpha\beta} \bar{\nabla}_\alpha \tau^a \bar{\nabla}_\beta \lambda_a.$$

The integration of this expression over the spatial section of spacetime leads to

$$\begin{aligned} K[\tau, \lambda] &= 2\kappa \int_\Sigma d^2x \varepsilon^{0\alpha\beta} \bar{\nabla}_\alpha \tau^a \bar{\nabla}_\beta \lambda_a \\ &= 2\kappa \int_{\partial\Sigma} dS_\alpha \varepsilon^{0\alpha\beta} (\tau^a \partial_\beta \lambda_a + \theta^a \bar{A}_{a\beta}), \end{aligned}$$

where we used the Stokes theorem and $\bar{F}^a_{\alpha\beta} = 0$. The second term in this expression is a functional of θ , hence it represents a trivial contribution to the central charge. Ignoring this term, we find that the essential piece of K is given by

$$K_{\text{ess}}[\tau, \lambda] = 2\kappa \int dS_\alpha \varepsilon^{0\alpha\beta} \tau^a \partial_\beta \lambda_a = 2\kappa \int_0^{2\pi} d\varphi \tau^a \partial_\varphi \lambda_a. \quad (5.4)$$

The same result follows from the surface integral expression (4.7b), as expected. At the canonical level, the related asymptotic symmetry is described by the affine (or Kac-Moody) extension of $sl(2, R)$ [8,9].

2. The second case is defined by a set of additional asymptotic conditions, which restrict the original asymptotic symmetry, as described bellow.

(a) First, we replace τ^a with the new gauge parameters ξ^μ , according to

$$\tau^a = \xi^\mu A^a_\mu.$$

The composition law is defined by $[\delta_\eta, \delta_\xi] \mathcal{A}_\mu^a = \delta_\chi \mathcal{A}_\mu^a$, with $\chi^\mu = \xi \cdot \partial \eta^\mu - \eta \cdot \partial \xi^\mu$. The new parameters ξ^μ correspond to diffeomorphisms, which is related to the gravitational content of the Chern-Simons theory. Note that, as a consequence of $A_+ = \mathcal{O}_1$, the parameter ξ^+ is effectively absent from the asymptotic region, $\xi^\mu A_\mu = \xi^r A_r + \xi^- A_- + \mathcal{O}_1$. As before, $\partial_+ A_\mu = \mathcal{O}_1$ and $\partial_+ \tau = \mathcal{O}_1$, which yields $\partial_+ \xi^\mu = \mathcal{O}_1$.

(b) In the second step, the asymptotic behaviour of A_μ fields is further restricted by the requirement

$$A_r = \alpha + \mathcal{O}_1,$$

where α is a constant element of the Lie algebra. Again, the field equations can be used to refine the asymptotics. This leads to

$$A_- = e^{-r\alpha}\Omega_-(x^-)e^{r\alpha} + \mathcal{O}_1,$$

with $\Omega_-(x^-)$ an arbitrary Lie algebra valued function of x^- . The new asymptotic conditions restrict the diffeomorphism parameters (ξ^r, ξ^-) to be independent of r , and $\tau \equiv \xi^\mu A_\mu$ takes the form $\tau = e^{-r\alpha}u(x^-)e^{r\alpha} + \mathcal{O}_1$. The conditions $A_+ = \mathcal{O}_1$ and $A_r = \alpha + \mathcal{O}_1$ are clearly invariant under the residual symmetry transformations of this form, while Ω_- is transformed as follows:

$$\delta\Omega_- = \partial_- u + [\Omega_-, u].$$

(c) Motivated by the relevant considerations in 3D gravity, we restrict (ξ^r, ξ^-) to be of the specific form [8,9]:

$$(\xi^r, \xi^-) = (\partial_\varphi \xi, \xi).$$

Such a choice leads to the conformal asymptotic symmetry of 3D gravity. It is straightforward to verify that the restricted parameters are consistent with the composition law of diffeomorphisms.

For the classical solution $\bar{A}$, which belongs to the set of asymptotic states described above, we choose

$$\bar{A}_+ = 0, \quad \bar{A}_r = \alpha, \quad \partial_+ \bar{A}_- = 0.$$

Using the change of parameters introduced in (a), one finds that the central charge of the restricted symmetry $K[\xi, \eta]$ is obtained from $K[\tau, \lambda]$ by the replacements $\tau^a \rightarrow \bar{\tau}^a \equiv \xi^\mu \bar{A}_\mu^a$ and $\lambda^a \rightarrow \bar{\lambda}^a \equiv \eta^\mu \bar{A}_\mu^a$:

$$K[\xi, \eta] = 2\kappa \int_{\partial\Sigma} dS_\alpha \varepsilon^{0\alpha\beta} (\bar{\tau}^a \partial_\beta \bar{\lambda}_a + \bar{\theta}^a \bar{A}_{a\beta}).$$

The property $\bar{A}_+ = 0$ implies the vanishing of the last term, since $\varepsilon^{abc} \bar{A}_{a\mu} \bar{A}_{b\nu} \bar{A}_{c\beta} = 0$. Then, one can derive an important identity, which immediately leads to

$$\begin{aligned} K[\xi, \eta] &= K_{\text{ess}}[\xi, \eta] + 2\kappa \int d\varphi \left(\alpha \cdot \bar{\Omega}_- \chi^r - \frac{1}{2} \bar{\Omega}_-^2 \chi^- \right), \\ K_{\text{ess}}[\xi, \eta] &\equiv 2\kappa \alpha^2 \int d\varphi \xi^r \partial_\varphi \eta^r. \end{aligned} \tag{5.5a}$$

The second term in $K[\xi, \eta]$ is a trivial piece of the central charge. Finally, replacing here the restricted gauge parameters as defined in (c), we obtain

$$K_{\text{ess}}[\xi, \eta] = 2\kappa \alpha^2 \int_0^{2\pi} d\varphi \partial_\varphi \xi \partial_\varphi^2 \eta. \tag{5.5b}$$

The structure of the asymptotic symmetry obtained in this way is described by the Virasoro algebra with classical central term, $c = -12 \times 4\pi\kappa\alpha^2$ [8,9].

If we change $\bar{A}_-$ and keep $\bar{A}_r = \alpha$ fixed, the change of the central charge is trivial. To understand the dependence of K_{ess} on α , it is useful to recall that $e^{r\alpha}$ defines the holonomy of the flat connection A_μ [11]. In the canonical framework, α defines the values of the conserved asymptotic generators. Thus, α is an essential characteristic of the chosen

asymptotic configuration, and consequently, changing α would mean a transition to another, non-equivalent asymptotic configuration, possessing a different central charge.

It is interesting to note that 3d gravity can be formulated in terms of two independent Chern-Simons gauge theories. In Riemannian or teleparallel theory we have $4\pi\kappa = \ell/8G$, $\alpha^2 = -1$, and $c_1 = c_2 = 3\ell/2G$, while in Riemann-Cartan gravity, one can have two different central charges [20].

B. Liouville theory

As our second example, we consider the Liouville theory on a cylinder. Using the coordinates $x^0 = t \in (-\infty, +\infty)$, $x^1 = \varphi \in [0, 2\pi)$, the Lagrangian and the related equations of motion read

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \phi)(\partial^\mu \phi) - \frac{\mu^2}{\alpha^2} e^{\alpha\phi}, \quad \square\phi + \frac{\mu^2}{\alpha} e^{\alpha\phi} = 0. \quad (5.6)$$

The transformation law

$$\delta_\varepsilon \phi = -\varepsilon^\mu \partial_\mu \phi - \frac{1}{\alpha} \partial_\mu \varepsilon^\mu \quad (5.7)$$

defines the symmetry of the theory, provided the parameters ε^μ satisfy the conditions

$$\partial_\mp \varepsilon^\pm = 0 \quad \Rightarrow \quad \varepsilon^\pm = \varepsilon^\pm(x^\pm),$$

where $x^\pm = x^0 \pm x^1$, and similarly for $\varepsilon^\pm$.

Let us now define the variable $\varphi \equiv \phi - \bar{\phi}$, where $\bar{\phi}$ is a fixed field configuration. Then, the Lagrangian takes the polynomial form (3.2) with the coefficients $V_i^\mu \equiv V^\mu$ and $V_i^{\mu\nu} \equiv V^{\mu\nu}$ given by

$$V^\mu = 0, \quad V^{\mu\nu} = \frac{1}{2} \eta^{\mu\nu}. \quad (5.8)$$

At the same time, the transformation law becomes

$$\delta_\varepsilon \varphi = \bar{R}(\varepsilon) - \varepsilon^\mu \partial_\mu \varphi, \quad \bar{R}(\varepsilon) \equiv -\varepsilon^\mu \partial_\mu \bar{\phi} - \frac{1}{\alpha} \partial_\mu \varepsilon^\mu. \quad (5.9)$$

The composition law of the group is obtained from the commutator algebra of these transformations: $[\delta_\eta, \delta_\varepsilon]\varphi = \bar{R}(\theta) - \theta^\mu \partial_\mu \varphi$, where $\theta^\mu \equiv \eta^\nu \partial_\nu \varepsilon^\mu - \varepsilon^\nu \partial_\nu \eta^\mu$.

Now, we are ready to calculate the central charge. Using the formula (3.5b), with the coefficients (5.8), we find

$$\mathcal{K}^\mu(\varepsilon, \eta) = \eta^{\mu\nu} \bar{R}(\varepsilon) \partial_\nu \bar{R}(\eta) - (\varepsilon \leftrightarrow \eta),$$

wherefrom

$$\begin{aligned} \mathcal{K}^0(\varepsilon, \eta) = & \left\{ \frac{1}{\alpha^2} \left[(\partial_+ \varepsilon^+) (\partial_+^2 \eta^+) + (\partial_- \varepsilon^-) (\partial_-^2 \eta^-) \right] \right. \\ & \left. + (\varepsilon^+ \eta^-) (\square \bar{\phi}) \partial_1 \bar{\phi} + \frac{1}{\alpha} (\square \bar{\phi}) \partial_1 (\varepsilon^+ \eta^-) \right\} - (\varepsilon \leftrightarrow \eta) + \partial_1 D + \mathcal{O}(\theta). \end{aligned} \quad (5.10)$$

In the above formula, the terms $\partial_1 D$ and $\mathcal{O}(\theta)$ are not explicitly displayed because they do not influence the central charge. The first is a spatial derivative of some quantity D , and consequently, vanishes after the integration over the compact spatial section of the cylinder. The second gives a trivial contribution to the central term since its dependence on ε and η is only through $\theta(\varepsilon, \eta)$.

We continue the analysis of the central charge by recalling that $\bar{\phi}$ must be a solution of the field equations. Using this fact, we find that the whole $\bar{\phi}$ dependent part of the equation (5.10) becomes a spatial derivative, and can be included in $\partial_1 D$. The final expression for the central charge is obtained by integrating $\mathcal{K}^0(\varepsilon, \eta)$ over the spatial section of the cylinder:

$$K_{\text{ess}}[\varepsilon, \eta] = \frac{1}{\alpha^2} \int_0^{2\pi} d\varphi \left[(\partial_+ \varepsilon^+) (\partial_+^2 \eta^+) + (\partial_- \varepsilon^-) (\partial_-^2 \eta^-) \right] - (\varepsilon \leftrightarrow \eta). \quad (5.11)$$

Two comments are in order: a) The obtained expression for central charge has no explicit $\bar{\phi}$ dependence. This is a consequence of the fact that $\bar{\phi}$ satisfies both conditions (a) and (b) given in section III A. b) The central charge (5.11) is nontrivial despite the fact that the spatial section of the cylinder has no boundary. This follows from our observation in section III B, that central terms are not always given as boundary integrals.

C. General relativity

In the tetrad formalism, general relativity with a cosmological constant is described by the Lagrangian density

$$\begin{aligned} \mathcal{L} &= -abR(A) - 2a\Lambda b \\ &= \frac{1}{2} a \varepsilon^{\mu\nu\lambda\rho} b^k{}_\lambda b^l{}_\rho \left(\partial_\mu A^{ij}{}_\nu + A^i{}_{s\mu} A^{sj}{}_\nu \right) - 2a\Lambda b, \end{aligned} \quad (5.12a)$$

where $b^i{}_\mu$ are tetrad fields, and $A^{ij}{}_\mu$ is the spin connection. Gauge symmetries of the theory are local translations and local Lorentz rotation, with parameters $\varepsilon^a = (\varepsilon^\mu, \varepsilon^{ij})$:

$$\begin{aligned} \delta_\varepsilon b^i{}_\mu &= \varepsilon^i{}_s b^s{}_\mu - (\partial_\mu \varepsilon^\lambda) b^i{}_\lambda - \varepsilon^\rho \partial_\rho b^i{}_\mu, \\ \delta_\varepsilon A^{ij}{}_\mu &= -\nabla_\mu \varepsilon^{ij} - (\partial_\mu \varepsilon^\lambda) A^{ij}{}_\lambda - \varepsilon^\rho \partial_\rho A^{ij}{}_\mu. \end{aligned} \quad (5.12b)$$

The commutator algebra $[\delta_\varepsilon, \delta_\tau] b^i{}_\mu = \delta_\theta b^i{}_\mu$ defines the composition law:

$$\begin{aligned} \theta^i{}_k &= \tau^i{}_m \varepsilon^m{}_k - \tau \cdot \partial \varepsilon^i{}_k - (\varepsilon \leftrightarrow \tau), \\ \theta^\rho &= \tau \cdot \partial \varepsilon^\rho - \varepsilon \cdot \partial \tau^\rho. \end{aligned}$$

After introducing

$$b^i{}_\mu = \bar{b}^i{}_\mu + \mathcal{B}^i{}_\mu, \quad A^{ij}{}_\mu = \bar{A}^{ij}{}_\mu + \mathcal{A}^{ij}{}_\mu,$$

where $(\bar{b}^i{}_\mu, \bar{A}^{ij}{}_\nu)$ is a solution of the field equations, we can expand $\mathcal{L}$ in powers of $(\mathcal{B}, \mathcal{A})$, leading to

$$V_{ij}^{\mu\nu} = 0, \quad V_{ij}^\mu (\partial_\mu \varphi^i) \varphi^j = \varepsilon^{\mu\nu\lambda\rho} \bar{b}^k{}_\lambda \mathcal{B}^l{}_\rho \partial_\mu \mathcal{A}^{ij}{}_\nu. \quad (5.13)$$

Then, equation (4.7b) yields directly the surface integral expression for the central charge. Indeed, we have

$$\begin{aligned} N^{0\beta}(\tau) &= V_{ij}^0 \varphi^i \bar{R}_a^{j\beta} \tau^a - V_{ij}^\beta \varphi^i \bar{R}_a^{j0} \tau^a \\ &= 2a \varepsilon_{ijkl}^{0\alpha\gamma\beta} \bar{b}^k{}_\gamma (-\tau^\rho \bar{b}^l{}_\rho) \mathcal{A}^{ij}{}_\alpha, \end{aligned}$$

and consequently,

$$\begin{aligned} K[\varepsilon, \tau] &= \int dS_\beta \left[\delta_\varepsilon N^{0\beta}(\tau) \right]_{\mathcal{A}, \mathcal{B}=0} \\ &= 2a \int dS_\beta \varepsilon_{ijkl}^{0\alpha\gamma\beta} \bar{b}^k{}_\gamma \bar{b}^l{}_\rho \left[\bar{\nabla}_\alpha \varepsilon^{ij} + (\partial_\alpha \varepsilon^\lambda) \bar{A}^{ij}{}_\lambda + \varepsilon \cdot \partial \bar{A}^{ij}{}_\alpha \right] \tau^\rho. \end{aligned} \quad (5.14)$$

This formula is the tetrad counterpart of the metric expression for central charge in general relativity [1,15].

VI. CONCLUDING REMARKS

In the canonical formalism, asymptotic symmetry is characterized by the PB algebra, which may have central extension. Motivated by this canonical structure, we carried out an investigation of the central term from the Lagrangian point of view. Our analysis is based on the assumption that the adopted asymptotic configuration of fields guarantees that the canonical generators are finite, differentiable and conserved. A simple canonical derivation of the central term leads to the basic formula (3.4). This result is then transformed into the Lagrangian definition of the central term (3.14), where the role of canonical charges is taken over by suitable normalized Noether charges. In the case of gauge theories, we found the general surface integral expression (4.7) for the central term. The results are then applied to several interesting examples, which are intended to illustrate the method. We expect that the present analysis of central term could be extended to a complete Lagrangian treatment of asymptotic symmetries, fully equivalent with the canonical structure (1.1).

ACKNOWLEDGMENTS

This work was supported by the Serbian Science Foundation, Serbia.

APPENDIX A: DIFFERENT FORMS OF THE NOETHER IDENTITY

In this appendix, we derive two alternative forms of the Noether identity.

After using the expansion (4.5) for F_i in (4.3a), the condition that the coefficients multiplying fields φ^i and their derivatives vanish, leads to

$$\begin{aligned} \bar{R}_a^i f_{ij} &= \partial_\rho (\bar{R}_a^{i\rho} f_{ij}) + \mathcal{O}_1, \\ \bar{R}_a^i f_{ij}^\lambda &= \bar{R}_a^{i\lambda} f_{ij} + \partial_\rho (\bar{R}_a^{i\rho} f_{ij}^\lambda) + \mathcal{O}_1, \\ \bar{R}_a^i f_{ij}^{(\mu\lambda)} &= \bar{R}_a^{i(\mu} f_{ij}^{\lambda)} + \partial_\rho (\bar{R}_a^{i\rho} f_{ij}^{(\mu\lambda)}) + \mathcal{O}_1, \\ 0 &= \sum_{\pi(\mu\lambda\rho)} \bar{R}_a^{i\mu} f_{ij}^{\lambda\rho}, \end{aligned} \quad (A1)$$

where the sum goes over all permutations π of (μ, λ, ρ) (the symmetrization).

The identity (3.10) is obtained from the invariance condition (3.7a). We now show that for gauge theories, this relation reduces to the form (A1). Indeed, replacing the expression (4.1) for $\bar{R}^i$ into (3.10), the arbitrariness of ε^a and $\partial\varepsilon^a$ leads to the relations

$$\begin{aligned} V_{ij}^\rho \partial_\rho \bar{R}_a^i + 2V_{ij} \bar{R}_a^i &= \partial_\mu (V_{ji}^\mu \bar{R}_a^i) + 2\partial_\mu (V_{ji}^{\mu\rho} \partial_\rho \bar{R}_a^i), \\ V_{ij}^\mu \bar{R}_a^i + V_{ij}^\lambda \partial_\lambda \bar{R}_a^{i\mu} + 2V_{ij} \bar{R}_a^{i\mu} &= V_{ji}^\mu \bar{R}_a^i + \partial_\lambda (V_{ji}^\lambda \bar{R}_a^{i\mu}) \\ &\quad + 2V_{ji}^{\mu\rho} \partial_\rho \bar{R}_a^i + 2\partial_\lambda (V_{ji}^{\lambda\mu} \bar{R}_a^i) + 2\partial_\lambda (V_{ji}^{\lambda\rho} \partial_\rho \bar{R}_a^{i\mu}), \\ 2V_{ij}^{(\rho} \bar{R}_a^{i\mu)} &= 2V_{ji}^{(\mu\rho)} \bar{R}_a^i + 2V_{ji}^{(\mu\lambda} \partial_\lambda \bar{R}_a^{i\rho)} + 2\partial_\lambda (V_{ji}^{\lambda(\rho} \bar{R}_a^{i\mu)}), \\ 0 &= \sum_{\pi(\rho\mu\lambda)} V_{ji}^{\rho\mu} \bar{R}_a^{j\lambda}, \end{aligned}$$

which can be rewritten in an equivalent form as

$$\begin{aligned} \bar{R}_a^i f_{ij} &= 2\partial_\mu (V_{ji}^\mu \bar{R}_a^i + V_{ji}^{\mu\rho} \partial_\rho \bar{R}_a^i), \\ \bar{R}_a^{i\mu} f_{ij} &= 2(V_{ji}^\mu \bar{R}_a^i + V_{ji}^{\mu\rho} \partial_\rho \bar{R}_a^i) \\ &\quad + 2\partial_\lambda (V_{ji}^\lambda \bar{R}_a^{i\mu} + V_{ji}^{\lambda\mu} \bar{R}_a^i + V_{ji}^{\lambda\rho} \partial_\rho \bar{R}_a^{i\mu}), \\ V_{ij}^{(\rho} \bar{R}_a^{i\mu)} &= V_{ij}^{(\rho\mu)} \bar{R}_a^i + V_{ji}^{(\mu\lambda} \partial_\lambda \bar{R}_a^{i\rho)} + \partial_\lambda (V_{ji}^{\lambda(\rho} \bar{R}_a^{i\mu)}), \\ 0 &= \sum_{\pi(\rho\mu\lambda)} V_{ji}^{\rho\mu} \bar{R}_a^{j\lambda}. \end{aligned} \tag{A2}$$

As one can verify directly, these relations coincide with the identities (A1).

APPENDIX B: THE NOETHER FORM OF $\mathcal{G}^\mu$

Here, we derive the formula (4.6) for the canonical current $\mathcal{G}^\mu$. Using the gauge transformations (4.1), the expression (3.5a) for $\mathcal{G}^\mu$ takes the form

$$\begin{aligned} \mathcal{G}^\mu &= -2 \left[V_{ij}^\mu \varphi^j + V_{ij}^{\mu\nu} \partial_\nu \varphi^j + \partial_\nu (\varphi^j V_{ji}^{\mu\nu}) \right] (\bar{R}_a^i \varepsilon^a + \bar{R}_a^{i\lambda} \partial_\lambda \varepsilon^a) + 2\partial_\nu (\varphi^j V_{ji}^{\mu\nu} \bar{R}^i) + \mathcal{O}_2 \\ &= -2 \left[V_{ij}^\mu \varphi^j + V_{ij}^{\mu\nu} \partial_\nu \varphi^j + \partial_\nu (\varphi^j V_{ji}^{\mu\nu}) \right] \bar{R}_a^i \varepsilon^a \\ &\quad + 2\partial_\lambda \left\{ \left[V_{ij}^\mu \varphi^j + V_{ij}^{\mu\nu} \partial_\nu \varphi^j + \partial_\nu (\varphi^j V_{ji}^{\mu\nu}) \right] \bar{R}_a^{i\lambda} \right\} \varepsilon^a \\ &\quad - 2\partial_\lambda \left\{ \left[V_{ij}^\mu \varphi^j + V_{ij}^{\mu\nu} \partial_\nu \varphi^j + \partial_\nu (\varphi^j V_{ji}^{\mu\nu}) \right] \bar{R}_a^{i\lambda} \varepsilon^a - \varphi^j V_{ji}^{\mu\lambda} \bar{R}^i \right\} + \mathcal{O}_2, \end{aligned}$$

This result can be rewritten as

$$\mathcal{G}^\mu = N_a^\mu \varepsilon^a + \partial_\rho N^{\mu\rho} + \mathcal{O}_2,$$

where

$$\begin{aligned} N_a^\mu &= 2 \left[(V_{ji}^\mu - \partial_\rho V_{ji}^{\mu\rho}) \bar{R}_a^i - \partial_\lambda ((V_{ji}^\mu - \partial_\rho V_{ji}^{\mu\rho}) \bar{R}_a^{i\lambda}) \right] \varphi^j \\ &\quad + 2 \left[-2V_{ij}^{(\mu\nu)} \bar{R}_a^i - (V_{ji}^\mu - \partial_\rho V_{ji}^{\mu\rho}) \bar{R}_a^{i\nu} + 2\partial_\lambda (V_{ij}^{(\mu\nu)} \bar{R}_a^{i\lambda}) \right] \partial_\nu \varphi^j \\ &\quad + 2 \left[2V_{ij}^{(\mu\nu)} \bar{R}_a^i \right] \partial_\nu \partial_\lambda \varphi^j + \mathcal{O}_2, \\ N^{\mu\lambda} &= 2 \left[(V_{ji}^\mu - \partial_\rho V_{ji}^{\mu\rho}) \varphi^j - 2V_{ij}^{(\mu\nu)} \partial_\nu \varphi^j \right] \bar{R}_a^{i\lambda} \varepsilon^a \\ &\quad + 2 \left[V_{ji}^{\mu\lambda} \varphi^j \right] (\bar{R}_a^i \varepsilon^a + \bar{R}_a^{i\rho} \partial_\rho \varepsilon^a) + \mathcal{O}_2. \end{aligned}$$

The form (4.5) of the field equations implies $N_a^\mu \varepsilon^a = F_i \bar{R}_a^{i\mu} \varepsilon^a + \mathcal{O}_2$, as we expected. Similarly, using the identities derived in Appendix A, one finds that the above expression for $N^{\mu\lambda}$ can be brought to the form (4.6b).

REFERENCES

- [1] J. D. Brown and M. Henneaux, Central charges in the canonical realization of asymptotic symmetries: an example from three-dimensional gravity, *Commun. Math. Phys.* **104** (1986) 207; see also M. Henneaux, Energy-momentum, angular momentum, and superscharge in 2+1 dimensions, *Phys. Rev.* **D29** (1984) 2766.
- [2] T. Regge and C. Teitelboim, Role of surface integrals in the Hamiltonian formulation of general relativity, *Ann. of Phys.* **88** (1974) 286;
- [3] L. Castellani, Symmetries of the constrained Hamiltonian systems, *Ann. Phys. (N. Y.)* **143** (1982) 357.
- [4] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, Princeton NJ, 1992); M. Blagojević, *Gravitation and Gauge Symmetries* (Institute of Physics Publishing, Bristol, 2002).
- [5] M. Henneaux, On the Poisson brackets of differentiable generators in classical field theory, *J. Math. Phys.* **27** (1986) 489.
- [6] M. Blagojević and M. Vasilíć, Asymptotic symmetries and conserved quantities in the Poincaré gauge theory, *Class. Quant. Grav.* **5** (1988) 1241; see also: Conservation laws in the teleparallel theory of gravity, *Phys. Rev. D* **64** (2001) 044010 [hep-th/0008159].
- [7] E. Witten, 2+1 dimensional gravity as an exactly soluble system, *Nucl. Phys.* **B311** (1988) 46; A. Achúcarro and P. Townsend, A Chern-Simons Action For Three-Dimensional Anti-De Sitter Supergravity Theories, *Phys. Lett.* **B180** (1986) 89.
- [8] M. Bañados, Global charges in Chern-Simons theory and 2+1 black hole, *Phys. Rev.* **D52** (1996) 5861 [hep-th/9405171]; M. Bañados, Three-dimensional quantum geometry and black holes, Invited talk at the Second Meeting “Trends in Theoretical Physics”, held in Buenos Aires, December, 1998, e-print hep-th/9901148; M. Bañados, Notes on black holes and three-dimensional gravity, Proceedings of the VIII Mexican School on Particles and Fields, AIP Conf. Proc. **490** (1999) 198 [hep-th/9903244].
- [9] M. Bañados, T. Brotz and M. Ortiz, Boundary dynamics and the statistical mechanics of the 2+1 dimensional black hole, *Nucl. Phys.* **B545** (1999) 340 [hep-th/9802076].
- [10] A. Strominger, Black hole entropy from near-horizon microstates, *JHEP* **9802** (1998) 009 [hep-th/9712251].
- [11] S. Carlip, Lectures on (2+1)-dimensional gravity, e-print gr-qc/9503024; S. Carlip, *Quantum Gravity in 2+1 Dimensions* (Cambridge University Press, Cambridge, 1998).
- [12] S. Carlip, Conformal field theory, (2 + 1)-dimensional gravity, and the BTZ Black Hole, e-print gr-qc/0503022.
- [13] I. M. Anderson and C. G. Torre, Asymptotic conservation laws in field theory, *Phys. Rev. Lett.* **77** (1996) 4109 [hep-th/9608008]; C. G. Torre, Local cohomology in field theory, e-print hep-th/9706092.
- [14] B. Julia and S. Silva, Currents and superpotentials in classical gauge invariant theories I. Local results with applications to perfect fluids and General relativity, *Class. Quant. Grav.* **15** (1998) 2173 [gr-qc/9804029]; S. Silva, On superpotentials and charge algebras of gauge theories, *Nucl. Phys.* **B558** (1999) 391 [hep-th/9809109]; B. Julia and S. Silva, Currents and superpotentials in classical gauge invariant theories II. Global aspects and the example of affine gravity, *Class. Quant. Grav.* **17** (2000) 4733 [gr-qc/0005127]; B. L. Julia, Holography of charges in gauge theories, e-print hep-th/0104231.

- [15] G. Barnich and F. Brandt, Covariant theory of asymptotic symmetries, conservation laws and central charges, e-print hep-th/0111246.
- [16] L. F. Abbot and S. Deser, Charge definition in nonabelian gauge theories, Phys. Lett. **B116** (1982) 259; Stability of gravity with a cosmological constant, Nucl. Phys. **B195** (1982) 76.
- [17] R. Wald and A. Zoupas, A General Definition of “Conserved Quantities” in General Relativity and Other Theories of Gravity [gr-qc/9911095].
- [18] E. Noether, Invariante Variationsprobleme, Nachrichten Kgl. Ges. d. Wiss. zu Goethingen, Math-Phys. Kl. **2** (1918) 235.
- [19] Y. Choquet-Bruhat, C. Morrette DeWitt and M. Dillard-Bleick, *Analysis, Manifolds and Physics* (North Holland, Amsterdam, 1989).
- [20] M. Blagojević and M. Vasilic, Asymptotic symmetries in 3d gravity with torsion, Phys. Rev. **D67** (2003) 084032 [gr-qc/0301051]; Asymptotic dynamics in 3d gravity with torsion, Phys. Rev. **D68** (2003) 124007 [gr-qc/0306070]; Three-dimensional gravity with torsion as a Chern-Simons gauge theory, Phys. Rev. **D68** (2003) 104023 [gr-qc/0307078].